

A photograph of an older man with white hair, smiling and looking to the side. He is wearing a red and black plaid shirt over a green knit scarf and has a large backpack on his back. The backpack contains a blue rolled-up mat and other gear. The background is a vast, flat, sandy landscape under a bright sky.

**Exceptional outcomes.  
*Proven again.***

**Japan IDE Clinical Study Results: Patency in complex SFA lesions.**

- 88%** 12-month primary patency in long, complex SFA lesions (n = 103)<sup>1</sup>
- 21.8 cm average lesion length
  - 65.7% chronic total occlusions (CTOs)
  - 84.5% TASC II C&D lesions



[goremedical.com/ap/viabahn](http://goremedical.com/ap/viabahn)

1. Ohki T, Kichikawa K, Yokoi H, *et al.* Outcomes of the Japanese multicenter Viabahn trial of endovascular stent grafting for superficial femoral artery lesions. *Journal of Vascular Surgery* 2017;66(1):130-142.e1.

W. L. Gore & Associates, Inc. | Flagstaff, AZ 86004

Products listed may not be available in all markets.

GORE®, VIABAHN®, and designs are trademarks of W. L. Gore & Associates.  
© 2018 W. L. Gore & Associates, Inc. AX1674-EN1 DECEMBER 2018

